



Tamiflu® (oseltamivir phosphate) is a prescription medicine used to treat the flu (influenza) in people two weeks of age and older who have had flu symptoms for no more than two days. Tamiflu can also reduce the chance of getting the flu in people one year and older. Tamiflu does not prevent bacterial infections that may happen with the flu. **Tamiflu is not a substitute for an annual flu vaccination.** Patients must not take Tamiflu if they are allergic to oseltamivir phosphate or any of the ingredients in Tamiflu. See below for additional Important Safety Information.

Tamiflu is FDA
approved for use
against influenza

A&B*

*Tamiflu has been studied only in strains of influenza that were circulating at the time of the clinical studies conducted to support FDA approval

Tamiflu is the only prescription
oral antiviral medicine
approved to treat patients



**TWO WEEKS OF
AGE TO
THOSE 65+**

**ACCORDING TO THE CDC,
CLINICAL STUDIES OF VIRUSES CAUSING FLU
DURING THE LAST THREE SEASONS
(2012-13, 2013-14, 2014-15) HAVE SHOWN
LOW RATES OF
RESISTANCE TO TAMIFLU***

*These low rates of resistance do not imply that the use of Tamiflu will have a positive outcome for any particular patient.

References to CDC are provided for informational purposes only. CDC does not endorse private products, services, or enterprises.

**AVAILABLE IN CAPSULES
AND A LIQUID FORMULA
(ORAL SUSPENSION)**

Lessen your chance of
upset stomach by
taking Tamiflu
with food



**SEE A DOCTOR QUICKLY. TREATMENT SHOULD BEGIN
WITHIN 2 DAYS (48 HOURS) OF FLU SYMPTOM ONSET.**



**Tamiflu helps block the
virus' ability to replicate
in the body.**

**IN STUDIES,
TAMIFLU HELPED
SHORTEN THE TIME**



**ADULTS WERE
SICK BY
1.3 DAYS
(30%
FASTER)**

**CHILDREN
AGES 1-12
WERE SICK BY
1.5 DAYS (26%
FASTER)**



**BY ATTACKING THE
FLU VIRUS AT ITS
SOURCE**

IMPORTANT SAFETY INFORMATION

- If patients have an allergic reaction or a severe rash with Tamiflu, they must stop taking it and contact their doctor right away. This may be very serious
- People with the flu, particularly children and adolescents, may be at an increased risk of seizure, confusion, or abnormal behavior early during their illness
- Patients must let their doctor know if they are pregnant, nursing, have heart problems, breathing problems, a weakened immune system (immunocompromised), kidney problems or other medical conditions as Tamiflu may not be right for them
- Patients must also tell their doctor about any medications they are taking or if they've received a nasal-spray flu vaccine in the past two weeks
- The most common side effects are mild to moderate nausea, vomiting, diarrhea and stomach pain

Report side effects to the FDA at (800) FDA-1088 or www.fda.gov/medwatch. Patients and caregivers may also report side effects to Genentech at (888) 835-2555. For additional important safety information, please visit www.tamiflu.com.