

## Patient Information

**EVERYSDI® [ev-RIZ-dee]  
(risdiplam)  
for oral solution**

**EVERYSDI® [ev-RIZ-dee]  
(risdiplam)  
tablets**

### What is EVERYSDI?

- EVERYSDI is a prescription medicine used to treat spinal muscular atrophy (SMA) in children and adults.

### Before taking EVERYSDI, tell your healthcare provider about all of your medical conditions, including if you:

- are pregnant or plan to become pregnant. If you are pregnant, or are planning to become pregnant, ask your healthcare provider for advice before taking this medicine. EVERYSDI may harm your unborn baby.
- are a woman who can become pregnant:
  - Before you start your treatment with EVERYSDI, your healthcare provider may test you for pregnancy. Because EVERYSDI may harm your unborn baby, you and your healthcare provider will decide if taking EVERYSDI is right for you during this time.
  - Talk to your healthcare provider about birth control methods that may be right for you. Use birth control while on treatment and for at least 1 month after stopping EVERYSDI.
  - **Pregnancy Registry.** There is a pregnancy registry for women who take EVERYSDI during pregnancy. The purpose of this registry is to collect information about the health of the pregnant woman and her baby. If you are pregnant or become pregnant while receiving EVERYSDI, tell your healthcare provider right away. Talk to your healthcare provider about registering with the EVERYSDI Pregnancy Registry. Your healthcare provider can enroll you in this registry or you can enroll by calling 1-833-760-1098 or visiting <https://www.evrysdipregnancyregistry.com>.
- are an adult male planning to have children: EVERYSDI may affect a man's ability to have children (fertility). If this is of concern to you, make sure to ask a healthcare provider for advice.
- are breastfeeding or plan to breastfeed. It is not known if EVERYSDI passes into breast milk and may harm your baby. If you plan to breastfeed, discuss with your healthcare provider about the best way to feed your baby while on treatment with EVERYSDI.

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Keep a list of them to show your healthcare provider, including your pharmacist, when you get a new medicine.

### How should I take EVERYSDI?

- For infants and children, your healthcare provider will determine the daily dose of EVERYSDI needed based on your child's age and weight. For adults, take 5 mg of EVERYSDI daily.
- Your healthcare provider will either prescribe:  
EVERYSDI for oral solution  
**Or**  
EVERYSDI tablet
- Your healthcare provider will tell you how long you or your child needs to take EVERYSDI. Do not stop treatment with EVERYSDI unless your healthcare provider tells you to.
  - Take EVERYSDI exactly as your healthcare provider tells you to take it. Do not change the dose without talking to your healthcare provider.
- Avoid getting EVERYSDI on your skin or in your eyes. If it gets on your skin, wash the area with soap and water. If it gets in your eyes, rinse your eyes with water.

### If you are taking EVERYSDI for oral solution:

- **see the detailed Instructions for Use that comes with it for information on how to take or give EVERYSDI for oral solution**
- You should receive EVERYSDI from the pharmacy as a liquid that can be given by mouth or through a feeding tube. The liquid solution is prepared by your pharmacist or other healthcare provider. If the medicine in the bottle is a powder, **do not use it**. Contact your pharmacist for a replacement.
- Take EVERYSDI one time daily with or without a meal at about the same time each day. Drink water afterwards to make sure EVERYSDI has been completely swallowed.
- In infants who are breastfed, EVERYSDI can be given before or after breastfeeding.
- Do not mix EVERYSDI with formula or milk.
- If you are unable to swallow and have a nasogastric (NG-tube) or gastrostomy tube (G-tube), EVERYSDI for oral solution can be given through the tube.

### Reusable Oral Syringes for EVRYSDI for Oral Solution

- Your pharmacist will provide you with the reusable oral syringes that are needed for taking your medicine and explain how to use them. You should receive 1 or 2 identical oral syringes depending on your prescribed daily dose.
- From the bottle, draw up (measure) the dose of EVRYSDI with these syringes, as they are made to protect the medicine from light.
- Take EVRYSDI right away after it has been drawn into the syringe. Do not store the EVRYSDI solution in the syringe. If EVRYSDI is not taken within 5 minutes of when it is drawn up, throw away the solution by pressing the plunger and prepare a new dose with the same syringe.
- Do not throw the syringes away because they are reusable.
- Wash the syringes per instructions after use.
- Contact your healthcare provider or pharmacist if your oral syringes are lost or damaged.

### If you are taking EVRYSDI Tablets:

- Take EVRYSDI one time daily with or without a meal at about the same time each day.
- Do not chew, cut, or crush the tablet.
- Swallow the EVRYSDI tablet whole with some water.

### Or

- You can take an EVRYSDI tablet by mouth after it is mixed with a teaspoon (5 mL) of room temperature non-chlorinated drinking water (e.g., filtered water). See the instructions below for taking EVRYSDI this way.

### EVRYSDI Tablets Mixed with Non-chlorinated Drinking Water (e.g., Filtered Water)

- Wash your hands before and after preparing or taking EVRYSDI tablets.
- Put EVRYSDI tablets in a small cup with a teaspoon (5 mL) of room temperature non-chlorinated drinking water (e.g., filtered water).
- Do not mix EVRYSDI tablets with any liquids other than non-chlorinated drinking water.
- Do not expose the EVRYSDI tablet mixture to sunlight.
- Swirl the cup gently until the tablet and non-chlorinated drinking water are fully mixed (though some particles may remain). This may take up to 3 minutes.
- Take EVRYSDI tablet mixture right away after mixing with non-chlorinated drinking water. If you do not take it within 10 minutes of adding non-chlorinated drinking water, throw the mixture away and make a new dose.
- After taking EVRYSDI tablet mixture, to make sure there is no medicine left, refill the small cup with at least a tablespoon (15 mL) of non-chlorinated drinking water, swirl, and take right away.
- If the EVRYSDI tablet mixture is spilled, use a dry paper towel to dry the area and then clean with soap and water. Throw the paper towel away in the trash and wash your hands with soap and water.
- Do not give the EVRYSDI tablet mixture via a nasogastric (NG-tube) or gastrostomy tube (G-tube).

### If you miss a dose of EVRYSDI:

- If you remember the missed dose within 6 hours of when you normally take EVRYSDI, then take the dose. Continue taking EVRYSDI at your usual time the next day.
- If you remember the missed dose more than 6 hours after you normally take EVRYSDI, skip the missed dose. Take your next dose at your usual time the next day.
- If you do not fully swallow the dose, or you vomit after taking a dose, **do not take** another dose of EVRYSDI to make up for that dose. Wait until the next day to take the next dose at your usual time.

### What are the possible side effects of EVRYSDI?

#### The most common side effects of EVRYSDI include:

- **For later-onset SMA:**
  - fever
  - diarrhea
  - rash
- **For infantile-onset SMA:**
  - fever
  - runny nose, sneezing, and sore throat (upper respiratory infection)
  - constipation
  - diarrhea
  - lung infection (lower respiratory infection)
  - vomiting
  - rash
  - cough

These are not all of the possible side effects of EVRYSDI. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

### **How should I store EVRYSDI?**

#### **EVRYSDI for Oral Solution:**

- Store EVRYSDI in the refrigerator between 36°F to 46°F (2°C to 8°C). Do not freeze.
- If necessary, EVRYSDI can be kept at room temperature up to 104°F (up to 40°C) for a combined total of 5 days. EVRYSDI can be removed from, and returned to, a refrigerator. The total combined time out of refrigeration should not be more than 5 days.
- Keep EVRYSDI in an upright position in the original amber bottle to protect from light.
- Throw away (discard) any unused portion of EVRYSDI 64 days after it is mixed by the pharmacist (constitution) or if EVRYSDI has been kept at room temperature (below 104°F [40°C]) for more than a total combined time of 5 days. Discard EVRYSDI if it has been kept above 104°F (40°C). Please see the Discard After date written on the bottle label. (See the **Instructions for Use** that comes with EVRYSDI for oral solution).

#### **EVRYSDI Tablets:**

- Store at room temperature between 68°F to 77°F (20°C to 25°C).
- Keep the bottle tightly closed in order to protect from moisture.

**Keep EVRYSDI, all medicines and syringes out of the reach of children.**

### **General information about the safe and effective use of EVRYSDI.**

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use EVRYSDI for a condition for which it was not prescribed. Do not give EVRYSDI to other people, even if they have the same symptoms you have. It may harm them. You can ask your pharmacist or healthcare provider for information about EVRYSDI that is written for health professionals.

### **What are the ingredients in EVRYSDI?**

**Active ingredient:** risdiplam

**Inactive ingredients:**

**EVRYSDI for Oral Solution:** ascorbic acid, disodium edetate dihydrate, isomalt, mannitol, polyethylene glycol 6000, sodium benzoate, strawberry flavor, sucralose, and tartaric acid.

**EVRYSDI Tablets:** colloidal silicon dioxide, crospovidone, mannitol, microcrystalline cellulose, polyethylene glycol 3350, polyvinyl alcohol, sodium stearyl fumarate, strawberry flavor, talc, tartaric acid, titanium dioxide, and yellow iron oxide.

Distributed by: **Genentech, Inc.**, A Member of the Roche Group, 1 DNA Way, South San Francisco, CA 94080-4990

EVRYSDI is a registered trademark of Genentech, Inc.

©2025 Genentech, Inc. All rights reserved.

For more information, go to [www.EVRYSDI.com](http://www.EVRYSDI.com) or call 1-833-387-9734.